



# Aloxi® (palonosetron) (Intravenous)

**Document Number: IC-0008** 

Last Review Date: 03/31/2023 Date of Origin: 10/17/2008

Dates Reviewed: 06/2009, 12/2009, 09/2010, 12/2010, 02/2011, 03/2011, 06/2011, 09/2011, 12/2011, 03/2012, 06/2012, 09/2012, 12/2012, 03/2013, 06/2013, 09/2013, 12/2013, 03/2014, 06/2014, 09/2014, 12/2014, 03/2015, 05/2015, 08/2015, 11/2015, 02/2016, 05/2016, 08/2016, 11/2016, 02/2017, 05/2017, 08/2017, 11/2017, 02/2018, 05/2018, 04/2019, 04/2020, 04/2021, 04/2022, 04/2023

## I. Length of Authorization

Coverage will be provided for 6 months and may be renewed, unless otherwise specified.

PONV: Coverage will be provided for 1 dose and may not be renewed.

## **II.** Dosing Limits

- A. Quantity Limit (max daily dose) [NDC unit]:
  - Aloxi 0.25 mg/5 mL solution for injection: 1 vial per 7 day supply
  - Aloxi 0.075 mg/1.5 mL solution for injection: 1 vial
- B. Max Units (per dose and over time) [HCPCS Unit]:

#### CINV:

• 10 billable units per 7 days

#### PONV:

• 3 billable units x 1 dose only

#### III. Initial Approval Criteria

Coverage is provided in the following conditions:

Prevention of Chemotherapy induced Nausea and vomiting (CINV) in Adults † 1-4,6

- Patient meets one of the following criteria:
  - o Patient is receiving highly emetogenic chemotherapy (HEC)\*; **OR**
  - Patient has failed\*\* with another 5HT3-antagonist (i.e., ondansetron or granisetron)
    while receiving the current chemotherapy regimen; AND
- Palonosetron is NOT covered for any of the following:



- o Breakthrough emesis
- o Repeat dosing in multi-day emetogenic chemotherapy regimens

# Prevention of Chemotherapy induced Nausea and vomiting (CINV) in Pediatric Patients † 1-4,6

- Patient is at least 1 month old and less than 17 years old; AND
- Patient is receiving emetogenic chemotherapy; AND
- Palonosetron is NOT covered for:
  - o Breakthrough emesis; OR
  - o Repeat dosing in multi-day emetogenic chemotherapy regimens

# Prevention of post-operative nausea and vomiting (PONV) in Adults † 1

# \*Highly emetogenic chemotherapy (HEC):

Highly Emetogenic Chemotherapy (HEC)					
Carboplatin	Carmustine	Cisplatin	Cyclophosphamide		
Dacarbazine	Doxorubicin	Epirubicin	Fam-trastuzumab deruxtecan-nxki		
Ifosfamide	Mechlorethamine	Melphalan ≥140 mg/m²	Sacituzumab govitecan- hziy		
Streptozocin					
The following can be considered HEC in certain patients					
Dactinomycin	Daunorubicin	Idarubicin	Irinotecan		
$\begin{array}{c} Methotrexate \\ \geq 250 mg/m^2 \end{array}$	Oxaliplatin	Trabectedin			
The following regimens can be considered HEC					
FOLFOX	FOLFIRI	FOLFIRINOX; FOLFOXIRI	AC (any anthracycline + cyclophosphamide)		

## \*\* Failure is defined as:

• Two or more documented episodes of vomiting attributed to the current chemotherapy regimen

† FDA Approved Indication(s); ‡ Compendium Recommended Indication(s); **\Phi** Orphan Drug

## IV. Renewal Criteria 1-3

Coverage may be renewed based upon the following criteria:



- Patient continues to meet indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; AND
- Disease response; AND
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: serotonin syndrome, severe QT prolongation, severe hypersensitivity reactions (including anaphylaxis and anaphylactic shock), etc.

# V. Dosage/Administration <sup>1</sup>

Indication	Dose	
Prevention of chemotherapy-induced nausea and vomiting in adults	Administer 0.25 mg intravenously, no more frequently than weekly, prior to emetogenic chemotherapy	
Prevention of chemotherapy-induced nausea and vomiting in pediatrics	Administer 20 mcg/kg (max of 1.5 mg) intravenously, no more frequently than weekly, prior to emetogenic chemotherapy	
Post-operative nausea and vomiting	Administer 0.075 mg intravenously immediately before the induction of anesthesia	

## VI. Billing Code/Availability Information

### **HCPCS Code**:

• J2469 – Injection, palonosetron HCl, 25 mcg: 1 billable unit = 25 mcg (0.025 mg)

#### NDC:

- Aloxi 0.25 mg/5 mL solution for injection; single-dose vial: 69639-103-xx\*
- Aloxi 0.075 mg/1.5 mL solution for injection; single-dose vial: 69639-103-xx (not commercially available)

#### VII. References

- 1. Aloxi [package insert]. Switzerland; Helsinn Healthcare SA; April 2020. Accessed March 2023.
- 2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) palonosetron. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc." To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2023.



<sup>\*</sup>Generics available from multiple manufacturers

- 3. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Antiemesis. Version 1.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc." To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2023.
- 4. Roila F, Molassiotis A, Herrstedt J, et al. MASCC and ESMO Consensus Guidelines for the Prevention of Chemotherapy and Radiotherapy-Induced Nausea and Vomiting: ESMO Clinical Practice Guidelines. Ann Oncol (2016) 27 (suppl 5): v119-v133.
- 5. Hesketh PJ, Kris MG, Basch E, et al. Antiemetics: American Society of Clinical Oncology Clinical Practice Guideline Update. J Clin Oncol. 2017 Oct 1;35(28):3240-3261.
- 6. Hesketh PJ, Kris MG, Basch E, et al. Antiemetics: ASCO Guideline Update. Journal of Clinical Oncology 2020 38:24, 2782-2797.

## Appendix 1 - Covered Diagnosis Codes

ICD-10	ICD-10 Description	
R11.0	Nausea	
R11.10	Vomiting, unspecified	
R11.11	Vomiting without nausea	
R11.12	Projectile vomiting	
R11.2	Nausea with vomiting, unspecified	
T41.0X5A	Adverse effect of inhaled anesthetics, initial encounter	
T41.1X5A	Adverse effect of intravenous anesthetics, initial encounter	
T41.205A	Adverse effect of unspecified general anesthetics, initial encounter	
T41.295A	Adverse effect of other general anesthetics, initial encounter	
T41.45XA	Adverse effect of unspecified anesthetic, initial encounter	
T45.1X5A	Adverse effect of antineoplastic and immunosuppressive drugs, initial encounter	
T45.1X5D	Adverse effect of antineoplastic and immunosuppressive drugs, subsequent encounter	
T45.1X5S	Adverse effect of antineoplastic and immunosuppressive drugs, sequela	
T45.95XA	Adverse effect of unspecified primarily systemic and hematological agent, initial encounter	
T45.95XD	Adverse effect of unspecified primarily systemic and hematological agent, subsequent encounter	
T45.95XS	Adverse effect of unspecified primarily systemic and hematological agent, sequela	
T50.905A	Adverse effect of unspecified drugs, medicaments and biological substances, initial encounter	
T50.905D	Adverse effect of unspecified drugs, medicaments and biological substances, subsequent	
T50.905S	Adverse effect of unspecified drugs, medicaments and biological substances, sequela	
T88.59XA	Other complications of anesthesia, initial encounter	
Z51.11	Encounter for antineoplastic chemotherapy	

ICD-10	ICD-10 Description
Z51.12	Encounter for antineoplastic immunotherapy

# Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs) and Local Coverage Article (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <a href="http://www.cms.gov/medicare-coverage-database/search.aspx">http://www.cms.gov/medicare-coverage-database/search.aspx</a>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions				
Jurisdiction	Applicable State/US Territory	Contractor		
E (1)	CA,HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC		
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC		
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)		
6	MN, WI, IL	National Government Services, Inc. (NGS)		
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.		
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)		
N (9)	FL, PR, VI	First Coast Service Options, Inc.		
J (10)	TN, GA, AL	Palmetto GBA, LLC		
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC		
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.		
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)		
15	КҮ, ОН	CGS Administrators, LLC		

