

## Perjeta® (pertuzumab) (Intravenous)

-E-

Document Number: MODA-0387

**Last Review Date: 10/03/2023**

**Date of Origin: 01/07/2019**

**Dates Reviewed: 01/2019, 04/2019, 07/2019, 10/2019, 01/2020, 04/2020, 07/2020, 10/2020, 01/2021, 05/2021, 07/2021, 10/2021, 02/2022, 05/2022, 07/2022, 10/2022, 01/2023, 04/2023, 7/2023, 10/2023**

### I. Length of Authorization <sup>1,2</sup>

Coverage is provided for 6 months and may be renewed (unless otherwise specified).

- Neoadjuvant and adjuvant treatment in Breast Cancer may be authorized up to a maximum of 1 year of treatment [18 cycles].

### II. Dosing Limits

#### A. Quantity Limit (max daily dose) [NDC Unit]:

Perjeta 420 mg/14mL solution for injection:

- **Loading Dose:** 2 vials
- **Maintenance Dose:** 1 vial every 21 days

#### B. Max Units (per dose and over time) [HCPCS Unit]:

- **Loading Dose:** 840 billable units x 1 dose
- **Maintenance Dose:** 420 billable units every 21 days

### III. Initial Approval Criteria <sup>1</sup>

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**

#### Universal Criteria <sup>1</sup>

- Left ventricular ejection fraction (LVEF) is within normal limits prior to initiating therapy and will be assessed at regular intervals (e.g., every 3 months) during treatment; **AND**
- Patient has human epidermal growth factor receptor 2 (HER2)-positive\* disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**
- Therapy will not be used in combination with pertuzumab/trastuzumab and hyaluronidase-zzxf (Phesgo); **AND**

### **Breast Cancer † ‡ 1-3,5-8,13,12e-15e,24e-26e**

- Used as neoadjuvant or preoperative therapy; **AND**
  - Patient has locally advanced, node positive, or inflammatory disease; **AND**
  - Used in combination with trastuzumab and a taxane-based regimen (e.g., docetaxel, paclitaxel, etc.); **OR**
- Used as adjuvant treatment; **AND**
  - Patient had node-positive disease at initial staging (i.e., cN+ or pN+); **AND**
    - Used in combination with trastuzumab and chemotherapy; **OR**
    - Used in combination with trastuzumab; **OR**
- Used for recurrent unresectable or metastatic disease; **AND**
  - Used as first-line therapy in combination with trastuzumab **AND** either paclitaxel or docetaxel; **OR**
  - Used as subsequent therapy in combination with trastuzumab with or without cytotoxic therapy †; **AND**
    - Patient was previously treated with trastuzumab and chemotherapy; **AND**
    - Patient has not previously received pertuzumab

### **Colorectal Cancer (CRC) † ‡ 2,9-12,16e**

- Used for RAS and BRAF wild-type (WT) disease in combination with trastuzumab; **AND**
- Used as subsequent therapy for progression of advanced or metastatic disease after at least one prior line of treatment in the advanced or metastatic disease setting; **AND**
- Patient has not previously received HER2-targeted therapy; **AND**
  - Used in one of the following:
    - Patient has mismatch repair proficient/microsatellite-stable (pMMR/MSS) disease; **OR**
    - Patient has mismatch repair deficient/microsatellite instability-high (dMMR/MSI-H) disease **AND** is not a candidate for or has progressed on checkpoint inhibitor immunotherapy

### **Head and Neck Cancers † 2,14,15,20e**

- Patient has salivary gland tumors; **AND**
- Used in combination with trastuzumab; **AND**
- Used for one of the following:
  - Recurrent disease with distant metastases
  - Unresectable locoregional recurrence with prior radiation therapy (RT)
  - Unresectable second primary with prior RT

## Biliary Tract Cancers (Gallbladder Cancer or Intra-/Extra-Hepatic Cholangiocarcinoma) ‡ 2,16,17

- Used as subsequent treatment for progression on or after systemic treatment for metastatic disease; **AND**
- Used in combination with trastuzumab

**Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.**

### \*HER2-positive overexpression criteria:

#### Breast, CNS, Head and Neck, and Biliary Tract Cancers: 3,4

- Immunohistochemistry (IHC) assay 3+; **OR**
- Dual-probe in situ hybridization (ISH) assay HER2/CEP17 ratio  $\geq 2.0$  AND average HER2 copy number  $\geq 4.0$  signals/cell; **OR**
- Dual-probe in situ hybridization (ISH) assay AND concurrent IHC indicating one of the following:
  - HER2/CEP17 ratio  $\geq 2.0$  AND average HER2 copy number  $< 4.0$  signals/cell AND concurrent IHC 3+; **OR**
  - HER2/CEP17 ratio  $< 2.0$  AND average HER2 copy number  $\geq 6.0$  signals/cell AND concurrent IHC 2+ or 3+; **OR**
  - HER2/CEP17 ratio  $< 2.0$  AND average HER2 copy number  $\geq 4.0$  and  $< 6.0$  signals/cell AND concurrent IHC 3+

#### Colorectal Cancer: 10,11

- Immunohistochemistry (IHC) assay 3+; **OR**
- Fluorescence in situ hybridization (FISH) HER2/CEP17 ratio  $\geq 2$  AND concurrent IHC 2+; **OR**
- Next-generation sequencing (NGS) panel HER2 amplification

❖ *If confirmed using an immunotherapy assay-<http://www.fda.gov/companiondiagnostics>*

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Ⓢ Orphan Drug

## IV. Renewal Criteria<sup>1</sup>

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**

- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: left ventricular dysfunction, severe infusion-related reactions, hypersensitivity reactions/anaphylaxis, etc.; **AND**
- Left ventricular ejection fraction (LVEF) obtained within the previous 3 months as follows:
  - Neoadjuvant and adjuvant treatment of breast cancer: LVEF is  $\geq 50\%$  OR LVEF has had an absolute decrease of  $< 10\%$  from baseline
  - All other indications: LVEF is  $> 45\%$  OR LVEF is 40% to 45% and absolute decrease is  $< 10\%$  from baseline

### Breast Cancer (neoadjuvant or adjuvant therapy) <sup>1,2</sup>

- Patient has not exceeded a maximum of 1 year of treatment (total of 18 cycles)

## V. Dosage/Administration <sup>1,10-13,15,16,18</sup>

Indication	Dose
Breast Cancer	Administer 840 mg intravenously x 1 dose, then 420 mg intravenously every 21 days thereafter until disease progression or unmanageable toxicity. <ul style="list-style-type: none"> <li>• Neoadjuvant therapy consists of 3 to 6 cycles prior to surgery.</li> <li>• Use for neoadjuvant and adjuvant treatment is limited to a total of 1 year of treatment (total of 18 cycles).</li> </ul> <p><i>*Note: When used for recurrent or metastatic breast cancer, therapy may be continued until disease progression or unmanageable toxicity.</i></p>
All other indications	Administer 840 mg intravenously x 1 dose, then 420 mg intravenously every 21 days thereafter until disease progression or unmanageable toxicity.

## VI. Billing Code/Availability Information

### HCPCS Code:

- J9306 – Injection, pertuzumab, 1 mg; 1 mg = 1 billable unit

### NDC:

- Perjeta 420 mg/14 mL solution for injection: 50242-0145-xx

## VII. References (STANDARD)

1. Perjeta [package insert]. South San Francisco, CA; Genentech, Inc.; February 2021. Accessed September 2023.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) pertuzumab. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL

COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed September 2023.

3. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Breast Cancer, Version 4.2023. National Comprehensive Cancer Network, 2023. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org. Accessed September 2023.
4. Wolff AC, Hammond EH, Allison KH, et al. Human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists Clinical Practice Guideline Focused Update. *J Clin Oncol* 2018;36:2105-2122.
5. Gianni L, Pienkowski T, Im YH, et al. 5-year analysis of neoadjuvant pertuzumab and trastuzumab in patients with locally advanced, inflammatory, or early-stage HER2-positive breast cancer (NeoSphere): a multicentre, open-label, phase 2 randomised trial. *Lancet Oncol*. 2016 Jun;17(6):791-800. doi: 10.1016/S1470-2045(16)00163-7. Epub 2016 May 11.
6. Baselga J, Cortes J, Kim SB, et al. CLEOPATRA Study Group. Pertuzumab plus trastuzumab plus docetaxel for metastatic breast cancer. *N Engl J Med*. 2012;366:109-119.
7. Schneeweiss A, Chia S, Hickish T, et al. Pertuzumab plus trastuzumab in combination with standard neoadjuvant anthracycline-containing and anthracycline-free chemotherapy regimens in patients with HER2-positive early breast cancer: a randomized phase II cardiac safety study (TRYPHAENA). *Ann Oncol*. 2013 Sep;24(9):2278-84.
8. von Minckwitz G, Procter M, de Azambuja E, et al. Adjuvant Pertuzumab and Trastuzumab in Early HER2-Positive Breast Cancer. *N Engl J Med*. 2017;377(2):122-131.
9. Hainsworth JD, Meric-Bernstam F, Swanton C, et al. Targeted Therapy for Advanced Solid Tumors on the Basis of Molecular Profiles: Results From MyPathway, an Open-Label, Phase IIa Multiple Basket Study. *Clin Oncol*. 2018 Feb 20;36(6):536-542.
10. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Colon Cancer, Version 2.2023. National Comprehensive Cancer Network, 2023. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org. Accessed September 2023.
11. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Rectal Cancer, Version 4.2023. National Comprehensive Cancer Network, 2023. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed September 2023.
12. Meric-Bernstam F, Hurwitz H, Raghav KPS, et al. Pertuzumab plus trastuzumab for HER2-amplified metastatic colorectal cancer (MyPathway): an updated report from a multicentre, open-label, phase 2a, multiple basket study. *Lancet Oncol*. 2019;20(4):518-530. doi:10.1016/S1470-2045(18)30904-5.

13. Swain SM, Ewer MS, Viale G, et al. Pertuzumab, trastuzumab, and standard anthracycline- and taxane-based chemotherapy for the neoadjuvant treatment of patients with HER2-positive localized breast cancer (BERENICE): a phase II, open-label, multicenter, multinational cardiac safety study. *Ann Oncol.* 2018;29(3):646-653. doi:10.1093/annonc/mdx773.
14. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Head and Neck Cancer, Version 2.2023. National Comprehensive Cancer Network, 2023. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed September 2023.
15. Kurzrock R, Bowles DW, Kang H, et al. Targeted therapy for advanced salivary gland carcinoma based on molecular profiling: results from MyPathway, a phase Iia multiple basket study. *Ann Oncol.* 2020;31(3):412-421.
16. Javle M, Borad MJ, Azad NS, et al. Pertuzumab and trastuzumab for HER2-positive, metastatic biliary tract cancer (MyPathway): a multicentre, open-label, phase 2a, multiple basket study. *Lancet Oncol.* 2021 Sep;22(9):1290-1300. Doi: 10.1016/S1470-2045(21)00336-3. Epub 2021 Jul 30.
17. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Biliary Tract Cancers, Version 2.2023. National Comprehensive Cancer Network, 2023. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed September 2023.
18. Lin NU, Pegram M, Sahebjam S, et al. Pertuzumab Plus High-Dose Trastuzumab in Patients With Progressive Brain Metastases and HER2-Positive Metastatic Breast Cancer: Primary Analysis of a Phase II Study. *J Clin Oncol.* 2021 Aug 20;39(24):2667-2675. doi: 10.1200/JCO.20.02822.

## VIII. References (ENHANCED)

- 1e. Baselga J, Cortes J, Kim SB, et al. Pertuzumab plus trastuzumab plus docetaxel for metastatic breast cancer. *N Engl J Med* 2012;366(2):109-119.
- 2e. Perez EA, Romond EH, Suman VJ, et al. Four-year follow-up of trastuzumab plus adjuvant chemotherapy for operable human epidermal growth factor receptor 2-positive breast cancer: joint analysis of data from NCCTG N9831 and NSABP B-31. *J Clin Oncol.* 2011 Sep 1;29(25):3366-73. doi: 10.1200/JCO.2011.35.0868. Epub 2011 Jul 18.
- 3e. Slamon D, Eiermann W, Robert N, et al. Adjuvant trastuzumab in HER2-positive breast cancer. *N Engl J Med.* 2011 Oct 6;365(14):1273-83. doi: 10.1056/NEJMoa0910383.
- 4e. Swain SM, Baselga J, Kim SB, et al. Pertuzumab, trastuzumab, and docetaxel in HER2-positive metastatic breast cancer. *N Engl J Med.* 2015 Feb 19;372(8):724-34. doi: 10.1056/NEJMoa1413513.

- 5e. Datko FM, D'Andrea G, Dickler M, et al. Phase II study of pertuzumab, trastuzumab, and weekly paclitaxel in patients with HER2-overexpressing metastatic breast cancer (MBC). *Journal of Clinical Oncology* 2012 30:27\_suppl, 134-134.
- 6e. Smyth LM, Iyengar NM, Chen MF, et al. Weekly paclitaxel with trastuzumab and pertuzumab in patients with HER2-overexpressing metastatic breast cancer: overall survival and updated progression-free survival results from a phase II study. *Breast Cancer Res Treat.* 2016 Jul;158(1):91-97. doi: 10.1007/s10549-016-3851-7. Epub 2016 Jun 15.
- 7e. Robert N, Leyland-Jones B, Asmar L, et al. Randomized phase III study of trastuzumab, paclitaxel, and carboplatin compared with trastuzumab and paclitaxel in women with HER2-overexpressing metastatic breast cancer. *J Clin Oncol.* 2006 Jun 20;24(18):2786-92.
- 8e. Andersson M, Lidbrink E, Bjerre K, et al. Phase III randomized study comparing docetaxel plus trastuzumab with vinorelbine plus trastuzumab as first-line therapy of metastatic or locally advanced human epidermal growth factor receptor 2-positive breast cancer: the HERNATA study. *J Clin Oncol.* 2011 Jan 20;29(3):264-71. doi: 10.1200/JCO.2010.30.8213. Epub 2010 Dec 13.
- 9e. Ellis PA, Barrios CH, Eiermann W, et al. Phase III, randomized study of trastuzumab emtansine (T-DM1) ± pertuzumab (P) vs trastuzumab + taxane (HT) for first-line treatment of HER2-positive MBC: Primary results from the MARIANNE study. *Journal of Clinical Oncology* 2015 33:15\_suppl, 507-507.
- 10e. Verma S, Miles D, Gianni L, et al. Trastuzumab emtansine for HER2-positive advanced breast cancer. *N Engl J Med.* 2012 Nov 8;367(19):1783-91. doi: 10.1056/NEJMoa1209124. Epub 2012 Oct 1.
- 11e. Krop IE, Kim SB, Martin AG, et al. Trastuzumab emtansine versus treatment of physician's choice in patients with previously treated HER2-positive metastatic breast cancer (TH3RESA): final overall survival results from a randomised open-label phase 3 trial. *Lancet Oncol.* 2017 Jun;18(6):743-754. doi: 10.1016/S1470-2045(17)30313-3. Epub 2017 May 16.
- 12e. Cameron D, Casey M, Oliva C, et al. Lapatinib plus capecitabine in women with HER2-positive advanced breast cancer: final survival analysis of a phase III randomized trial. *Oncologist.* 2010;15(9):924-34. doi: 10.1634/theoncologist.2009-0181. Epub 2010 Aug 24.
- 13e. Blackwell KL, Burstein HJ, Storniolo AM, et al. Overall survival benefit with lapatinib in combination with trastuzumab for patients with human epidermal growth factor receptor 2-positive metastatic breast cancer: final results from the EGF104900 Study. *J Clin Oncol.* 2012 Jul 20;30(21):2585-92. doi: 10.1200/JCO.2011.35.6725. Epub 2012 Jun 11.
- 14e. von Minckwitz G, du Bois A, Schmidt M, et al. Trastuzumab beyond progression in human epidermal growth factor receptor 2-positive advanced breast cancer: a german breast group 26/breast international group 03-05 study. *J Clin Oncol.* 2009 Apr 20;27(12):1999-2006. doi: 10.1200/JCO.2008.19.6618. Epub 2009 Mar 16.
- 15e. Baselga J, Gelmon KA, Verma S, et al. Phase II trial of pertuzumab and trastuzumab in patients with human epidermal growth factor receptor 2-positive metastatic breast cancer that progressed during prior trastuzumab therapy. *J Clin Oncol.* 2010 Mar 1;28(7):1138-44. doi: 10.1200/JCO.2009.24.2024. Epub 2010 Feb 1.
- 16e. Sartore-Bianchi A, Trusolino L, Martino C, et al. Dual-targeted therapy with trastuzumab and lapatinib in treatment-refractory, KRAS codon 12/13 wild-type, HER2-positive

- metastatic colorectal cancer (HERACLES): a proof-of-concept, multicentre, open-label, phase 2 trial. *Lancet Oncol.* 2016 Jun;17(6):738-746. doi: 10.1016/S1470-2045(16)00150-9.
- 17e. Murthy RK, Loi S, Okines A, et al. Tucatinib, Trastuzumab, and Capecitabine for HER2-Positive Metastatic Breast Cancer [published correction appears in *N Engl J Med.* 2020 Feb 6;382(6):586]. *N Engl J Med.* 2020;382(7):597-609. doi:10.1056/NEJMoa1914609.
- 18e. Modi S, Saura C, Yamashita T, et al. Trastuzumab Deruxtecan in Previously Treated HER2-Positive Breast Cancer. *N Engl J Med.* 2020;382(7):610-621. doi:10.1056/NEJMoa1914510.
- 19e. Thorpe L, Schrock A, Erlich R, et al. Significant and durable clinical benefit from trastuzumab in 2 patients with HER2-amplified salivary gland cancer and a review of the literature. *Head Neck* 2017 Mar;39(3):E40-E44. doi: 10.1002/hed.24634. Epub 2016 Dec 22.
- 20e. Takahashi H, Tada Y, Saotome T, et al. Phase II Trial of Trastuzumab and Docetaxel in Patients With Human Epidermal Growth Factor Receptor 2-Positive Salivary Duct Carcinoma. *J Clin Oncol* 2019 Jan 10;37(2):125-134. doi: 10.1200/JCO.2018.00545. Epub 2018 Nov 19.
- 21e. Jhaveri KL, Wang XV, Makker V, et al. Ado-trastuzumab emtansine (T-DM1) in patients with HER2-amplified tumors excluding breast and gastric/gastroesophageal junction (GEJ) adenocarcinomas: results from the NCI-MATCH trial (EAY131) subprotocol Q. *Ann Oncol.* 2019 Nov 1;30(11):1821-1830. doi: 10.1093/annonc/mdz291.
- 22e. Saura C, Oliveira M, Feng YH, et al. NALA Investigators. Neratinib Plus Capecitabine Versus Lapatinib Plus Capecitabine in HER2-Positive Metastatic Breast Cancer Previously Treated With  $\geq 2$  HER2-Directed Regimens: Phase III NALA Trial. *J Clin Oncol.* 2020 Sep 20;38(27):3138-3149. doi: 10.1200/JCO.20.00147. Epub 2020 Jul 17.
- 23e. Rugo HS, Im SA, Cardoso F, et al. SOPHIA Study Group. Efficacy of Margetuximab vs Trastuzumab in Patients With Pretreated ERBB2-Positive Advanced Breast Cancer: A Phase 3 Randomized Clinical Trial. *JAMA Oncol.* 2021 Apr 1;7(4):573-584. doi: 10.1001/jamaoncol.2020.7932.
- 24e. Taghian A. and Merajver S. (Oct 2022). Inflammatory breast cancer: Clinical features and treatment. In Hayes DF, Pierce LJ, Chagpar AB, et al. (Eds.), *UptoDate*. Accessed September 2023. Available from: [https://www.uptodate.com/contents/inflammatory-breast-cancer-clinical-features-and-treatment?search=inflammatory%20breast%20cancer&source=search\\_result&selectedTitle=1~48&usage\\_type=default&display\\_rank=1#H4025803376](https://www.uptodate.com/contents/inflammatory-breast-cancer-clinical-features-and-treatment?search=inflammatory%20breast%20cancer&source=search_result&selectedTitle=1~48&usage_type=default&display_rank=1#H4025803376).
- 25e. Sikov W.M. (July 2023). Neoadjuvant therapy for patients with HER2-positive breast cancer. In Burstein HJ, Vora SR (Eds.), *UptoDate*. Accessed September 2023. Available from: [https://www.uptodate.com/contents/neoadjuvant-therapy-for-patients-with-her2-positive-breast-cancer?sectionName=TIMING%20OF%20HER2-DIRECTED%20AGENTS&search=inflammatory%20breast%20cancer%20treatment&topicRef=768&anchor=H1845125338&source=see\\_link#H1845125338](https://www.uptodate.com/contents/neoadjuvant-therapy-for-patients-with-her2-positive-breast-cancer?sectionName=TIMING%20OF%20HER2-DIRECTED%20AGENTS&search=inflammatory%20breast%20cancer%20treatment&topicRef=768&anchor=H1845125338&source=see_link#H1845125338).
- 26e. Burstein HJ (Apr 2023). Adjuvant systemic therapy for HER2-positive breast cancer. In Isaacs C and Vora SR. (Eds.), *UptoDate*. Accessed September 2023. Available from: <https://www.uptodate.com/contents/adjuvant-systemic-therapy-for-her2-positive-breast-cancer?sectionName=Non-anthracycline>



based%20therapy&search=inflammatory%20breast%20cancer&topicRef=106774&anchor=H1352264107&source=see\_link#H1237051481.

- 27e. Siena S, Di Bartolomeo M, Raghav KPS, et al. A phase II, multicenter, open-label study of trastuzumab deruxtecan in patients with HER2-expressing metastatic colorectal cancer (mCRC): DESTINY-CRC01. *J Clin Oncol* 2020;38(suppl; abstr 4000).
- 28e. Bartsch R, Wenzel C, Altorjai G, et al. Capecitabine and Trastuzumab in Heavily Pretreated Metastatic Breast Cancer. *J Clin Oncol* 2007;25:3853-3858.
- 29e. Lee YR, Huh SJ, Lee DH, et al. Phase II Study of Vinorelbine Plus Trastuzumab in HER2 Overexpressing Metastatic Breast Cancer Pretreated with Anthracyclines and Taxanes. *J Breast Cancer*. 2011;14(2):140-146. doi:10.4048/jbc.2011.14.2.140.
- 30e. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Central Nervous System Cancers, Version 1.2023. National Comprehensive Cancer Network, 2023. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed September 2023.
- 31e. Strickler JH, Cercek A, Siena S, et al. Tucatinib plus trastuzumab for chemotherapy-refractory, HER2-positive, RAS wild-type unresectable or metastatic colorectal cancer (MOUNTAINEER): a multicentre, open-label, phase 2 study. *Lancet Oncol*. 2023 May; 24(5): 496-508. doi.org/10.1016/S1470-2045(23)00150-X.
- 32e. Robson M, Im SA, Senkus E, et al. Olaparib for metastatic breast cancer in patients with a germline BRCA mutation. *N Engl J Med* 2017;377:523-533.
- 33e. Litton JK, Rugo HS, Ettl J, et al. Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation. *N Engl J Med* 2018;379:753-763.
- 34e. Piccart M, Procter M, Fumagalli D, et al. APHINITY Steering Committee and Investigators. Adjuvant Pertuzumab and Trastuzumab in Early HER2-Positive Breast Cancer in the APHINITY Trial: 6 Years' Follow-Up. *J Clin Oncol*. 2021 May 1;39(13):1448-1457. doi: 10.1200/JCO.20.01204.
- 35e. von Minckwitz G, Huang CS, Mano MS, et al. Trastuzumab Emtansine for Residual Invasive HER2-Positive Breast Cancer. *N Engl J Med*. 2019 Feb 14;380(7):617-628.
- 36e. Magellan Health, Magellan Rx Management. Perjeta Clinical Literature Review Analysis. Last updated September 2023. Accessed September 2023.

## Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C06.9	Malignant neoplasm of mouth, unspecified
C07	Malignant neoplasm of parotid gland
C08.0	Malignant neoplasm of submandibular gland
C08.1	Malignant neoplasm of sublingual gland
C08.9	Malignant neoplasm of major salivary gland, unspecified
C18.0	Malignant neoplasm of cecum

ICD-10	ICD-10 Description
C18.2	Malignant neoplasm of ascending colon
C18.3	Malignant neoplasm of hepatic flexure
C18.4	Malignant neoplasm of transverse colon
C18.5	Malignant neoplasm of splenic flexure
C18.6	Malignant neoplasm of descending colon
C18.7	Malignant neoplasm of sigmoid colon
C18.8	Malignant neoplasm of overlapping sites of large intestines
C18.9	Malignant neoplasm of colon, unspecified
C19	Malignant neoplasm of rectosigmoid junction
C20	Malignant neoplasm of rectum
C21.8	Malignant neoplasm of overlapping sites of rectum, anus and anal canal
C22.1	Intrahepatic bile duct carcinoma
C23	Malignant neoplasm of gallbladder
C24.0	Malignant neoplasm of extrahepatic bile duct
C24.8	Malignant neoplasm of overlapping sites of biliary tract
C24.9	Malignant neoplasm of biliary tract, unspecified
C50.011	Malignant neoplasm of nipple and areola, right female breast
C50.012	Malignant neoplasm of nipple and areola, left female breast
C50.019	Malignant neoplasm of nipple and areola, unspecified female breast
C50.021	Malignant neoplasm of nipple and areola, right male breast
C50.022	Malignant neoplasm of nipple and areola, left male breast
C50.029	Malignant neoplasm of nipple and areola, unspecified male breast
C50.111	Malignant neoplasm of central portion of right female breast
C50.112	Malignant neoplasm of central portion of left female breast
C50.119	Malignant neoplasm of central portion of unspecified female breast
C50.121	Malignant neoplasm of central portion of right male breast
C50.122	Malignant neoplasm of central portion of left male breast
C50.129	Malignant neoplasm of central portion of unspecified male breast
C50.211	Malignant neoplasm of upper-inner quadrant of right female breast
C50.212	Malignant neoplasm of upper-inner quadrant of left female breast
C50.219	Malignant neoplasm of upper-inner quadrant of unspecified female breast
C50.221	Malignant neoplasm of upper-inner quadrant of right male breast
C50.222	Malignant neoplasm of upper-inner quadrant of left male breast
C50.229	Malignant neoplasm of upper-inner quadrant of unspecified male breast
C50.311	Malignant neoplasm of lower-inner quadrant of right female breast
C50.312	Malignant neoplasm of lower-inner quadrant of left female breast
C50.319	Malignant neoplasm of lower-inner quadrant of unspecified female breast

ICD-10	ICD-10 Description
C50.321	Malignant neoplasm of lower-inner quadrant of right male breast
C50.322	Malignant neoplasm of lower-inner quadrant of left male breast
C50.329	Malignant neoplasm of lower-inner quadrant of unspecified male breast
C50.411	Malignant neoplasm of upper-outer quadrant of right female breast
C50.412	Malignant neoplasm of upper-outer quadrant of left female breast
C50.419	Malignant neoplasm of upper-outer quadrant of unspecified female breast
C50.421	Malignant neoplasm of upper-outer quadrant of right male breast
C50.422	Malignant neoplasm of upper-outer quadrant of left male breast
C50.429	Malignant neoplasm of upper-outer quadrant of unspecified male breast
C50.511	Malignant neoplasm of lower-outer quadrant of right female breast
C50.512	Malignant neoplasm of lower-outer quadrant of left female breast
C50.519	Malignant neoplasm of lower-outer quadrant of unspecified female breast
C50.521	Malignant neoplasm of lower-outer quadrant of right male breast
C50.522	Malignant neoplasm of lower-outer quadrant of left male breast
C50.529	Malignant neoplasm of lower-outer quadrant of unspecified male breast
C50.611	Malignant neoplasm of axillary tail of right female breast
C50.612	Malignant neoplasm of axillary tail of left female breast
C50.619	Malignant neoplasm of axillary tail of unspecified female breast
C50.621	Malignant neoplasm of axillary tail of right male breast
C50.622	Malignant neoplasm of axillary tail of left male breast
C50.629	Malignant neoplasm of axillary tail of unspecified male breast
C50.811	Malignant neoplasm of overlapping sites of right female breast
C50.812	Malignant neoplasm of overlapping sites of left female breast
C50.819	Malignant neoplasm of overlapping sites of unspecified female breast
C50.821	Malignant neoplasm of overlapping sites of right male breast
C50.822	Malignant neoplasm of overlapping sites of left male breast
C50.829	Malignant neoplasm of overlapping sites of unspecified male breast
C50.911	Malignant neoplasm of unspecified site of right female breast
C50.912	Malignant neoplasm of unspecified site of left female breast
C50.919	Malignant neoplasm of unspecified site of unspecified female breast
C50.921	Malignant neoplasm of unspecified site of right male breast
C50.922	Malignant neoplasm of unspecified site of left male breast
C50.929	Malignant neoplasm of unspecified site of unspecified male breast
C78.00	Secondary malignant neoplasm of unspecified lung
C78.01	Secondary malignant neoplasm of right lung
C78.02	Secondary malignant neoplasm of left lung
C78.6	Secondary malignant neoplasm of retroperitoneum and peritoneum

ICD-10	ICD-10 Description
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
Z85.038	Personal history of other malignant neoplasm of large intestine

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC