

## Signifor® LAR (pasireotide) (Intramuscular)

Document Number: IC-0228

Last Review Date: 11/04/2025

Date of Origin: 01/06/2015

Dates Reviewed: 10/2015, 10/2016, 08/2018, 08/2019, 08/2020, 08/2021, 08/2022, 08/2023, 11/2024, 11/2025

### I. Length of Authorization

- Initial: Prior authorization validity will be provided initially for 6 months.
- Renewal: Prior authorization validity may be renewed every 12 months thereafter.

### II. Dosing Limits

**Max Units (per dose and over time) [HCPCS Unit]:**

- **Acromegaly**
  - 60 billable units (60 mg) every 28 days
- **Cushing's disease**
  - 40 billable units (40 mg) every 28 days

### III. Initial Approval Criteria <sup>1</sup>

Prior authorization validity is provided in the following conditions:

- Patient is at least 18 years of age; **AND**
- Confirmation that fasting plasma glucose, hemoglobin A1c, liver enzyme tests, electrocardiogram (ECG), serum magnesium and serum potassium have been evaluated prior to starting treatment; **AND**
- Patients with diabetes mellitus have stable glycemic control and are on optimized anti-diabetic treatment prior to the start of therapy; **AND**

#### Universal Criteria <sup>1</sup>

- Patient does not have severe hepatic impairment (i.e., Child-Pugh Class C); **AND**
- Patient has not received a long-acting somatostatin analog within the last 4 weeks; **AND**

#### Acromegaly † Φ <sup>1,4,5,10,14</sup>

- Patient's diagnosis is confirmed by one of the following:
  - Unequivocally elevated (age-adjusted) serum insulin-like growth factor-1 (IGF-1); **OR**

- Equivocally elevated (age-adjusted) serum IGF-1 AND inadequate suppression of growth hormone (GH) after a glucose load; **AND**
- Patient has documented inadequate response to surgery and/or radiotherapy or it is not an option for the patient; **AND**
- Baseline GH and IGF-1 blood levels have been obtained (renewal will require reporting of current levels); **AND**
- Will not be used in combination with oral octreotide or with GH-analog (e.g., pegvisomant)

#### **Cushing's Disease † ☐<sup>1,9,12,13</sup>**

- Confirmed diagnosis of endogenous Cushing's disease in which the patient's hypercortisolism is not a result of chronic administration of high-dose glucocorticoids or other physiologic conditions; **AND**
- Treatment of patient's disease with pituitary surgery has not been curative OR the patient is not a candidate for pituitary surgery; **AND**
- Baseline 24-hour urinary free cortisol (UFC) level, Adrenocorticotrophic hormone (ACTH), late-night salivary cortisol (LNSC), and/or serum cortisol level have been obtained (renewal will require reporting of current levels)

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); ☐ Orphan Drug

## **IV. Renewal Criteria<sup>1</sup>**

Prior authorization validity can be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: uncontrolled hyperglycemia, diabetes, ketoacidosis, bradycardia, QT prolongation, liver test elevations (e.g., alanine aminotransferase [ALT] or aspartate aminotransferase [AST]), cholelithiasis (gallstones) and complications of cholelithiasis (e.g., cholecystitis or cholangitis), pituitary hormone (e.g., thyroid, adrenal, gonadal) deficiencies/severe adrenal insufficiency, malabsorption of dietary fats (steatorrhea, stool discoloration, loose stools, etc.), etc.; **AND**

#### **Acromegaly<sup>1,4,5,10</sup>**

- Disease response as indicated by an improvement in signs and symptoms compared to baseline; **AND**
  - Reduction of growth hormone (GH) from pre-treatment baseline; **OR**
  - Age-adjusted normalization of serum IGF-1

## Cushing's Disease <sup>1,9,12,13</sup>

- Disease response indicated by reduction in urinary free cortisol (UFC), plasma adrenocorticotrophic hormone (ACTH), late-night salivary cortisol (LNSC), and/or serum cortisol levels from baseline

## V. Dosage/Administration <sup>1</sup>

Indication	Dose
Acromegaly	Initiate at 40 mg administered by intramuscular injection once every 4 weeks (28 days). <ul style="list-style-type: none"><li>– Titrate dosage based on treatment response and tolerability up to maximum 60 mg every 4 weeks for patients who have not normalized GH and/or IGF-1 levels after 3 months of treatment with the 40 mg dose.</li></ul>
Cushing's Disease	Initiate at 10 mg administered by intramuscular injection once every 4 weeks (28 days). <ul style="list-style-type: none"><li>– Titrate dosage based on treatment response and tolerability up to maximum 40 mg every 4 weeks for patients who have not normalized 24-hour urinary free cortisol (UFC) after 4 months of treatment with the 10 mg dose.</li></ul>

## VI. Billing Code/Availability Information

### HCPCS Code:

- J2502 - Injection, pasireotide long acting, 1 mg; 1 billable unit = 1 mg

### NDC(s):

- Signifor LAR 10 mg single-use kit: 55292-0139-xx
- Signifor LAR 20 mg single-use kit: 55292-0140-xx
- Signifor LAR 30 mg single-use kit: 55292-0141-xx
- Signifor LAR 40 mg single-use kit: 55292-0142-xx
- Signifor LAR 60 mg single-use kit: 55292-0143-xx

## VII. References

1. Signifor LAR [package insert]. Bridgewater, NJ; Recordati Rare Diseases, Inc.; July 2024. Accessed October 2025.
2. Colao A, Bronstein MD, Freda P, et al. Pasireotide versus octreotide in acromegaly: a head-to-head superiority study. J Clin Endocrinol Metab. 2014 Mar; 99(3):791-9. doi: 10.1210/jc.2013-2480. Epub 2014 Jan 13.
3. Gadelha MR, Bronstein MD, Brue T, et al. Pasireotide versus continued treatment with octreotide or lanreotide in patients with inadequately controlled acromegaly (PAOLA): a

- randomised, phase 3 trial. *Lancet Diabetes Endocrinol.* 2014 Nov; 2(11):875-84. doi: 10.1016/S2213-8587(14)70169-X. Epub 2014 Sep 24.
4. Giustina A, Chanson P, Kleinberg D, et al. Expert consensus document: A consensus on the medical treatment of acromegaly. *Nat Rev Endocrinol.* 2014 Apr; 10(4):243-8. doi: 10.1038/nrendo.2014.21. Epub 2014 Feb 25.
  5. Katznelson L, Laws ER Jr, Melmed S, et al. Acromegaly: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab.* 2014 Nov; 99(11):3933-51. doi: 10.1210/jc.2014-2700. Epub 2014 Oct 30.
  6. Lacroix A, Gu F, Gallardo W, et al; Pasireotide G2304 Study Group. Efficacy and safety of once-monthly pasireotide in Cushing's disease: a 12 month clinical trial. *Lancet Diabetes Endocrinol.* 2018 Jan;6(1):17-26. doi: 10.1016/S2213-8587(17)30326-1. Epub 2017 Oct 12. Erratum in: *Lancet Diabetes Endocrinol.* 2018 Jan;6(1):e1.
  7. Biller BM, Grossman AB, Stewart PM, et al. Treatment of Adrenocorticotropin-Dependent Cushing's Syndrome: A Consensus Statement. *J Clin Endocrinol Metab.* July 2008, 93(7):2454-2462.
  8. Hur KY, Kim JH, Kim BJ, et al. Clinical Guidelines for the Diagnosis and Treatment of Cushing's Disease in Korea. *Endocrinol Metab (Seoul).* 2015 Mar; 30(1): 7–18.
  9. Nieman LK, Biller BMK, Findling JW, et al. Treatment of Cushing's Syndrome: An Endocrine Society Clinical Practice Guideline. *The Journal of Clinical Endocrinology & Metabolism*, Volume 100, Issue 8, 1 August 2015, Pages 2807–2831.
  10. Fleseriu M, Biller BMK, Freda PU, et al. A Pituitary Society update to acromegaly management guidelines. *Pituitary* 24, 1–13 (2021). <https://doi.org/10.1007/s11102-020-01091-7>
  11. Giustina A, Barkhoudarian G, Beckers A et al. Multidisciplinary management of acromegaly: A consensus. *Rev Endocr Metab Disord* 21, 667–678 (2020). <https://doi.org/10.1007/s11154-020-09588-z>
  12. Fleseriu M, Auchus R, Bancos I, et al. Consensus on diagnosis and management of Cushing's disease: a guideline update. *Lancet Diabetes Endocrinol* 2021; 9: 847–75. Published Online October 20, 2021. [https://doi.org/10.1016/S2213-8587\(21\)00235-7](https://doi.org/10.1016/S2213-8587(21)00235-7)
  13. Price J, Pivonello R, Tabarin A, et al. Use of late-night salivary cortisol to monitor response to medical treatment in Cushing's disease. *Eur J Endocrinol.* 2020 Feb;182(2):207-217. doi: 10.1530/EJE-19-0695.
  14. Melmed S, Katznelson L (Apr 2023). Treatment of acromegaly. In Snyder PJ, Martin KA (Eds.) *UpToDate*. Accessed October 06, 2025. Available from: <https://www.uptodate.com/contents/treatment-of-acromegaly>

## Appendix A – Non-Quantitative Treatment Limitations (NQTL) Factor Checklist

Non-quantitative treatment limitations (NQTLs) refer to the methods, guidelines, standards of evidence, or other conditions that can restrict how long or to what extent benefits are provided under a health plan. These may include things like utilization review or prior authorization. The utilization management NQTL applies comparably, and not more stringently, to mental health/substance use disorder (MH/SUD) Medical Benefit Prescription Drugs and medical/surgical (M/S) Medical Benefit Prescription Drugs. The table below lists the factors that were considered in designing and applying prior authorization to this drug/drug group, and a summary of the conclusions that Prime's assessment led to for each.

Factor	Conclusion
Indication	Yes: Consider for PA
Safety and efficacy	No: PA not a priority
Potential for misuse/abuse	No: PA not a priority
Cost of drug	Yes: Consider for PA

## Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
E22.0	Acromegaly and pituitary gigantism
E24.0	Pituitary-dependent Cushing's disease
E34.4	Constitutional tall stature

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCA/LCD): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC